Information about changes to medicines or treatments on the NHS: Changes to immediate-release fentanyl prescribing

The NHS will be asking doctors to stop or greatly reduce the prescribing of some medicines from December 2017. This is because the medicines are:

- Not as safe as other medicines OR
- Not as good (effective) as other medicines OR
- More expensive than other medicines that do the same thing.

One of these medicines is immediate-release fentanyl.

This document will explain why the changes are happening and where you can get more information and support.

What is fentanyl?

Fentanyl is a strong painkiller. It is available as an immediate-release painkiller in various dosage forms: tablets, lozenges, films and nasal spray. "Immediate-release" means that it acts very quickly, and it is sometimes used for the treatment of "breakthrough" pain in adults with cancer who are already receiving other, longer-acting strong painkillers.

It is also used for other types of pain, but it does not have a licence* for these and can be addictive.

*Having a licence means that a medicine has passed tests to ensure that it is effective, safe and manufactured to appropriate quality standards.

Why does the NHS want to reduce prescribing of immediaterelease fentanyl?

The National Institute for Health and Care Excellence (NICE) says that immediate-release fentanyl should not be the first choice for "breakthrough" pain in adults with cancer – it should only be used if other options haven't worked. This is likely to be for just a small number of people.

Also, the amount of immediate-release fentanyl being prescribed nationally makes it likely that in many cases it is being used for other types of pain than cancer. Immediate-release fentanyl can cause addiction, so we want to reduce use in these cases.

This **does not** apply to slow-release versions of fentanyl, which come in patch form. Fentanyl patches will still be available on prescription.

Will my immediate-release fentanyl be stopped?

If you are taking immediate-release fentanyl for cancer pain it will not be stopped.

If you are taking immediate-release fentanyl for other types of pain it will be reviewed by your doctor. You may be supported to stop taking immediate-release fentanyl and offered a different painkiller if needed.

Where can I find more information and support?

- You can speak to your local pharmacist, GP or the person who prescribed the medication to you
- National and local charities can offer advice and support, for example:
 - Pain Concern: 0300 123 0789 http://painconcern.org.uk/
 - Pain UK: https://painuk.org
 - Cancer Research: www.cancerresearchuk.org/
 - MacMillan: www.macmillan.org.uk/information-and-support/coping/side-effects-and-symptoms/pain
- The NICE information for patients can be found at: www.nice.org.uk/guidance/cg140/ifp/chapter/About-this-information
- The Patients Association can also offer support and advice: www.patients-association.org.uk/ or call 020 8423 8999

Find out more about the medicines that are being stopped or reduced: https://www.england.nhs.uk/medicines/items-which-should-not-be-routinely-prescribed/

Find out more about licensed and unlicensed medicines:

https://www.gov.uk/drug-safety-update/off-label-or-unlicensed-use-of-medicines-prescribers-responsibilities#a-licensed-medicine-meets-acceptable-standards-of-efficacy-safety-and-quality

If you have any questions about fentanyl prescribing please email them to: england.medicines@nhs.net